

WHY ARE THERE DWARF SPHEROIDAL GALAXIES?

RENYUE CEN¹

Draft version February 1, 2008

ABSTRACT

There appears to exist a conflict between the standard structure formation theory and the existence of the faint dwarf spheroidal galaxies in the Local Group. Theoretical considerations suggest that a cold dark matter universe has been a very hostile place for the formation of small galaxies. In particular, gas would not have been able to cool and condense to form stars inside dark matter halos with a velocity dispersion ≤ 10 km/s since before the cosmological reionization epoch. In other words, one should not expect to see any major star formation activities in dwarf galaxies with a velocity dispersion ≤ 10 km/s during the past 12 Gyr, according to the current structure formation theory. The real universe, on the other hand, shows that all dwarf spheroidal galaxies in the Local Group have velocity dispersions ≤ 10 km/s and extended and recent star formation activities are quite common in them.

This apparent conflict between cold dark matter models and local observations can be resolved, if one allows the cold dark matter particles to decay relatively recently. The resolution comes about in that the dwarf spheroidal galaxies with a velocity dispersion of ~ 10 km/s seen today were born, in this picture, in a small fraction of halos with a velocity dispersion of ~ 20 km/s, where gas can be retained and is able to cool and contract to form stars. The presently observed lower velocity dispersion (~ 10 km/s) of these dwarf spheroidal galaxies is a consequence of the decay of one half of the dark matter in and subsequent expansion of the halos since redshift $z \sim 2$.

Subject headings: Cosmology: theory – galaxies: evolution – galaxies: formation – galaxies: halos – Local Group

1. INTRODUCTION

In the standard picture of structure formation, over the redshift interval $z \sim 10 - 100$ a sufficient amount of molecular hydrogen may be able to form and enable gas to cool and condense at temperature $T \ll 10^4$ K, capable of assembling stellar systems with a velocity dispersion $\ll 10$ km/s (Haiman, Thoul, & Loeb 1996; Tegmark et al. 1997). However, a very small fraction of total baryons collapsed in these first generation stellar systems would have formed enough stars to generate enough sub-13.6eV photons (to which the universe is largely transparent even then) to completely photo-dissociate hydrogen molecules well before enough hydrogen ionizing photons (> 13.6 eV) have been produced to reionize the universe.

Subsequently, gas cooling and hence collapsing is not possible in halos with a virial temperature $T_v < 10^4$ K (or baryonic masses $< 10^8[(1 + z_v)/10]^{-3/2}$; Haiman, Rees, & Loeb 1997). However, an early soft X-ray background could enhance the formation of hydrogen molecules (by providing extra free electrons) thus may somewhat prolong the period of gas cooling in halos with a virial temperature $T_v < 10^4$ K (Haiman, Abel, & Rees 2000). Once hydrogen molecules are destroyed, star formation becomes inactive (there may be exceptional regions enriched with metals where metal cooling at $T < 10^4$ is possible thus some star formation activities may take place there), until the nonlinear scale of the density perturbations reaches a virial temperature $T \sim 10^4$ K.

At that point a sufficient number of galaxies in halos of virial temperature $T \geq 10^4$ K (with hydrogen collisional excitation cooling being the primary coolant) quickly form enough stars and the universe gets reionized at a very rapid pace (Gnedin & Ostriker 1997). Between the epoch of

molecular hydrogen elimination and the completion of cosmological reionization some gas may passively sit in halos with $T_v \leq 10^4$ K but is incapable of cooling and condensing to form stars. Unfortunately, even this already settled gas will be driven out of the shallow halo potential wells (on a timescale of sound crossing time $\ll 1$ Gyr), when the universe is reionized, leaving the halos virtually gasless (Barkana & Loeb 1999) and depriving them of chance for any further star formation.

After reionization, the temperature of the intergalactic medium of the universe is raised to $\geq 10^4$ K (Miralda-Escudé & Rees 1994) and subsequent accretion of gas onto halos with $T_v \leq 10^4$ K is prohibited (Efstathiou 1992; Thoul & Weinberg 1996; Quinn, Katz, & Efstathiou 1996; Kepner, Babul, & Spergel 1997; Navarro & Steinmetz 1997; Kitayama & Ikeuchi 2000; Gnedin 2000b). Reionization of once-ionized helium at lower redshift (perhaps at $z \sim 3$) may still increase the temperature of the intergalactic medium (Hui & Gnedin 1997; Schaye et al. 2000), while additional shock heating due to breaking of long density waves progressively raises the temperature of the intergalactic medium further at somewhat lower redshift (Cen & Ostriker 1999).

In summary, it does not seem to be possible to (bottom-up) form isolated stellar systems with a velocity dispersion < 10 km/s since redshift at least as high as $z = 6$, at which the universe seems to have already been reionized (Fan et al. 2000). The age of the universe then is ≤ 1 Gyr. In other words, star formation in dwarf galaxies with a velocity dispersion ≤ 10 km/s can only occur ≥ 12 Gyr ago (assuming that the present age of the universe is 13 Gyr) and no later episodes of star formation are allowed. This is a generic feature of cold dark matter models as well as

¹Princeton University Observatory, Princeton University, Princeton, NJ 08544; cen@astro.princeton.edu

warm or collisional dark matter models.

However, observations of our local neighborhood, the Local Group, show that there exist a dozen or so dwarf spheroidal galaxies, all of which have a velocity dispersion ≤ 10 km/s (8.5 ± 2.0 km/s seems to bracket the velocity dispersions of all the Local Group dwarf spheroidal galaxies; van den Bergh 2000). These dwarf spheroidal galaxies have complicated star formation histories but it is clear that extended and recent star formation activities are common. For example, the star formation in Fornax may be as recent as < 1 Gyr ago (Saviane, Held, & Bertelli 2000), so are Sculptor (Monkiewicz et al. 1999) and Sagittarius (Bellazzini, Ferraro, & Buonanno 1999); Carina experienced three significant episodes of star formation at ~ 15 , ~ 7 and ~ 3 Gyr ago, with the main burst of star formation during the episode 7 Gyr ago (Hurley-Keller, Mateo, & Nemec 1998); In Leo I 70-80% of stars formed between 7 and 1 Gyr ago (Gallart et al. 1999).

How can one reconcile this clear theoretical prediction with solid local observations? One possible solution is that these dwarf spheroidal galaxies are produced during the formation of or interactions (such as close encounters or mergers) between giant galaxies. However, this explanation seems unlikely because the central dark matter density of the observed dwarf spheroidal galaxies appears to be higher than that of giants; fragments of larger galaxies are unlikely to have a higher density than their parents (Kormendy & Freeman 1998).

In this *Letter*, we suggest that this apparent conflict may be resolved in the decaying cold dark matter model (ADCDM) recently proposed (Cen 2000). This is not because gas can be accreted onto halos with a velocity dispersion < 10 km/s in ADCDM rather it is because halos seen today with a velocity dispersion 10 km/s have a velocity dispersion 20 km/s at birth in ADCDM, onto which gas accretion is not prohibited or already settled gas would not have been driven out of the halos in the cosmological reionization phase.

2. FORMATION OF DWARF SPHEROIDAL GALAXIES IN DECAYING COLD DARK MODEL

Dwarf spheroidal galaxies have very high mass-to-light ratios (van den Bergh 2000, Fig 15.3; Hirashita, Tamura, & Takeuchi 2000) and thus are dark matter dominated (Faber & Lin 1983; Lin & Faber 1983). Observations indicate that the stellar mass function in them is very similar to globular clusters in giant galaxies down to very small stellar mass, implying that it is unlikely that the dark matter in the dwarf spheroidal galaxies may be stellar, since globular clusters do not contain a significant amount of dark matter (Feltzing, Gilmore, & Wyse 1999). This then brings out our assumption that dwarf spheroidal galaxies are cold dark matter dominated stellar systems on all scales in the context of cold dark matter models. Therefore, the dynamics of dwarf spheroidal galaxies is mostly determined by the cold dark matter within; dynamically, baryons evolve passively inside dark matter halos of dwarf spheroidal galaxies.

The decaying cold dark matter model (Cen 2000) was recently suggested to alleviate the small-scale power excess problem at $z = 0$ in the canonical cosmological constant dominated cold dark matter model (Ostriker & Steinhardt 1995), which has been remarkably successful on larger

scales. In this model one half of the cold dark matter particles decay into relativistic particles by zero redshift with a decay time of $1.4t_0$ (t_0 is the current age of the universe). By $z = 2$ about 10% of cold dark matter particles have decayed. Hence, in this model, virialized halos are not static but evolve at low redshift ($z < 2$) when a substantial amount of dark matter starts to decay. Because the dark matter decay time scale is much longer than the dynamic time of a halo, a halo primarily experiences an adiabatic expansion when the cold dark matter particles decay at low redshift. Since the baryons do not play a significant dynamic role, the dynamics of dark matter halos is particularly simple. Basically, an isolated halo will grow larger in size (radius) by a factor of approximately two and its total mass becomes smaller by a factor of approximately two, after one half of the dark matter has decayed. Therefore, the velocity dispersion of such a halo is approximately reduced by a factor of two from $z \sim 2$ to $z = 0$.

As a consequence, a halo seen today with a velocity dispersion of 10 km/s had a velocity dispersion of 20 km/s at birth and had maintained this velocity dispersion until $z \sim 2$ in ADCDM. Unlike halos with a velocity dispersion of 10 km/s, where accretion of photoionized gas is completely suppressed, gas accretion onto halos with a velocity dispersion of 20 km/s is permitted. Besides, throughout the cosmological reionization phase these halos would have retained gas that was settled in before cosmological reionization. This may resolve the apparent conflict between the existence of extended star formation activities in dwarf spheroidal galaxies and the cold dark matter model.

3. DISCUSSION

Although star formation is, in principle, not prohibited in halos with a velocity dispersion of 20 km/s and could occur over an extended period throughout the life of the universe, gas accretion onto them may be significantly hindered by photoionization heating and other heating processes. In fact, simulations have shown that photoionization heating can cause suppression of gas accretion onto halos with a velocity dispersions up to $50 - 70$ km/s (Thoul & Weinberg 1996; Quinn, Katz, & Efstathiou 1996; Navarro & Steinmetz 1997; Gnedin 2000b). But detailed hydrodynamic simulations with sufficient mass and spatial resolutions to resolve the halos in question are required to quantify this effect definitively.

However, even if gas accretion onto halos with a velocity dispersion of 20 km/s is completely suppressed after the cosmological reionization epoch, these halos may still have enough gas that was accreted at earlier times; these halos could have accreted gas before the cosmological reionization epoch and retained the gas, unlike halos with a velocity dispersion of ≤ 10 km/s (Barkana & Loeb 1999). It is relevant to discuss the effect of supernovae from starbursts in dwarf spheroidal galaxies. Mac Low & Ferrara (1999) have shown that mass ejection efficiency is very low for galaxies with mass $M > 10^7 M_\odot$. Since the dwarf spheroidal galaxies today have mass $M > 2 \times 10^7 M_\odot$ (and higher at high redshift in our model), it seems possible that they have retained most of their gas over their lifetime. This would then have permitted extended star formation activities in them, although star formation activities could be intermittent due to internal supernova heating (Burkert

& Ruiz-Lapuente 1997; Hirashita 2000); this is consistent with the observed star formation histories of local dwarf spheroidal galaxies.

It is worth noting that the observationally inferred velocity dispersion thus mass of the dwarf spheroidal galaxies is based on the assumption that these galaxies are in dynamical equilibrium. If they are not in dynamic equilibrium, then their mass would be lower and dark matter content may not be as high, which would exacerbate the problem considered here; i.e., without dark matter halos, the star formation activities of the dwarf spheroidal galaxies would be even more difficult to understand. Considerations of tidal perturbations of dwarf spheroidal galaxies by giants prefer the existence of a deep potential well of dark matter (Hirashita, Kamaya, & Takeuchi 1999); in other words, without dark matter halos, it will be very difficult to maintain the orderly existence of dwarf spheroidal galaxies in the vicinities of large galaxies as in the Local Group environment. Under the assumption of dynamical equilibrium one finds that the average density within the virial radius of the dwarf spheroidal galaxies is approximately $10^6 \rho_{crit}$ (where ρ_{crit} is the present critical density of the universe), indicating an early formation epoch ($z \geq 10$) of these halos (e.g., Peebles 1997). This early epoch of formation has the virtue of collecting gas easily before cosmological reionization and does not necessarily require substantial accretion of gas at later epochs, as discussed above. Using Press-Schechter (1974) theory one estimates that the number density of halos with a velocity dispersion > 20 km/s is $10^3 - 10^4$ Mpc $^{-3}$ at $z = 10$ in the Λ CDM model, significantly higher than the number density of the Local Group dwarf spheroidal galaxies (~ 1 Mpc $^{-3}$). Clearly, a large number of halos with a velocity dispersion 20 km/s could have formed before cosmological reionization.

Klypin et al. (1999) have shown that the abundance of halos with a velocity dispersion of ≥ 20 km/s in the Λ CDM model at $z = 0$ is roughly a factor of 3 higher than the abundance of the observed dwarf spheroidal galaxies with a velocity dispersion of ~ 10 km/s in the Local Group. Assuming that the only effect of introducing decaying cold dark matter particles is to reduce the velocity dispersion of halos by a factor of two, the difference between their simulations and observations implies that only about 1/3 of halos with a velocity dispersion of ~ 20 km/s in the Λ CDM model that have not been cannibalized end up as dwarf spheroidal galaxies with a velocity dispersion of ~ 10 km/s as seen in the Local Group. The fate of the remaining 2/3 of dwarf spheroidal galaxy halos with a velocity dispersion of ≥ 10 km/s has interesting consequences. One possibility is that they may have been tidally disturbed and thus are more dispersed to be detected easily perhaps at stages later than the Sagittarius dwarf galaxy (Ibata, Gilmore, & Irwin 1994) or is already part of the halo stars of giants. The second possibility is that these dwarf spheroidal galaxies are just further away and they are not yet detected due to their low surface brightness. The third possibility is that they were neither tidally disrupted nor formed enough stars. In this case, it is unclear whether they are gas rich or not. Perhaps likely is that the gas content of these “dark” dwarf spheroidal galaxies is strongly environment dependent; those closer to the giants may suffer effects such as ram pressure stripping. It

is also possible that these “dark” dwarf galaxies may contain mostly ionized gas, thus their gas may not be easily detectable in HI 21 cm surveys (e.g., Young 2000).

Interestingly, there is an intriguing coincidence. In the vast majority of all the observed dwarf spheroidal galaxies the star formation rate declines sharply about 10 Gyr ago (Grebel 1998; Mateo 1998; Gnedin 2000a), which corresponds to redshift $z \sim 2$ in the Λ CDM model (Cen 2000). Incidentally, this is about the redshift after which cold dark matter particles start to decay significantly and hence dwarf galaxy halos start to expand significantly. Two effects could arise. First, the expansion of the halos cause a simultaneous expansion of the baryonic gas thus reduces the baryonic surface density. Second, the lowering of the halo velocity dispersion further reduces the accretion rate of photoionized gas onto the halos. Both effects reduce star formation rate and could have caused the sharp decline of star formation rate in the observed dwarf spheroidal galaxies ~ 10 Gyr ago.

Finally, the scenario presented here may be tested by future observations. It is predicted that there should be no dwarf spheroidal galaxies with a velocity dispersion of ≤ 10 km/s at $z \geq 2$; there should be a minimum velocity dispersion of approximately 20 km/s for dwarf spheroidal galaxies at $z \geq 2$. The dwarf spheroidal galaxies should be more compact at $z \geq 2$ than today; they should be smaller in size by a factor of ~ 2 and surface brightness higher by a factor of ~ 4 (perhaps more considering stellar evolution) at $z \geq 2$ than today. NGST may be able to detect such dwarf spheroidal galaxies at $z \sim 2$ (Stockman & Mather 1999).

4. CONCLUSIONS

A host of theoretical arguments indicate that no significant star formation activity should have taken place in dwarf spheroidal galaxies with a velocity dispersion ≤ 10 km/s in the past 12 Gyr, in any cold (or warm or collisional) dark matter universe. Observations, on the other hand, tell us that all dwarf spheroidal galaxies in the Local Group have a velocity dispersion ≤ 10 km/s and much more recent (than 12 Gyr ago) star formation activities in them are common. Therefore, we face a conflict between theory and observations.

We suggest that this apparent conflict may be resolved in the decaying cold dark matter model. In this model these dwarf spheroidal galaxies were born in halos with a velocity dispersion of ~ 20 km/s, in which gas may be retained and/or accreted, and cooling and condensation could occur for an extended period of time. Thus, extended and recent star formation activities in them would have been possible. The presently observed lower velocity dispersion (≤ 10 km/s) of these dwarf spheroidal galaxies results from the decay of one half of the dark matter in and subsequent expansion of the halos since redshift $z \sim 2$.

Future observations such as by NGST may be able to detect high redshift dwarf spheroidal galaxies. It is predicted by the current scenario that they should be smaller, have higher surface brightness and higher velocity dispersion at high redshift.

This research is supported in part by grants AST93-18185 and ASC97-40300. I thank Zoltan Haiman and Jerry Ostriker for many useful comments that significantly

improved the paper.

REFERENCES

- Barkana, R., & Loeb, A. 1999, *ApJ*, 523, 54
 Bellazzini, M., Ferraro, F.R., & Buonanno, R. 1999, *MNRAS*, 307, 619
 Burkert, A., & Ruiz-Lapuente, P. 1997, *ApJ*, 480, 297
 Cen, R. 2000, *ApJ*, in press, astro-ph/0005206
 Cen, R., & Ostriker, J.P. 1999, *ApJ*, 514, 1
 Da Costa, G.S., Armandroff, T.E., Caldwell, N., & Seitzer, P. 2000, *AJ*, 119, 705
 Efstathiou, G. 1992, *MNRAS*, 256, 43
 Faber, S.M., & Lin, D.N.C., 1983, *ApJ*, 266, L17
 Fan, X., et al. 2000, *AJ*, 120, 1167
 Feltzing, S., Gilmore, G., & Wyse, R.F.G. 1999, *ApJ*, 516, L17
 Gallart, C., Freedman, W.L., Aparicio, A., Bertelli, G., & Chiosi, C. 1999, *AJ*, 118, 2245
 Gnedin, N.Y. 2000a, *ApJ*, 535, L75
 Gnedin, N.Y. 2000b, *ApJ*, 542, 535
 Gnedin, N.Y., & Ostriker, J.P. 1997, *ApJ*, 486, 581
 Grebel, E. 1998, *IAUS*, 192, 1
 Haiman, Z., Thoul, A.A., & Loeb, A. 1996, *ApJ*, 464, 523
 Haiman, Z., Rees, M.J., & Loeb, A. 1997, *ApJ*, 476, 458
 Haiman, Z., Abel, T., & Rees, M.J. 2000, *ApJ*, 534, 11
 Hirashita, H. 2000, *PASP*, 52, 107
 Hirashita, H., Kamaya, H., & Takeuchi, T.T. 1999, *PASP*, 51, 375
 Hirashita, H., Tamura, N., & Takeuchi, T.T. 2000, "New Cosmological Data and the Values of the Fundamental Parameters", *IAUS* 201, p64 (Manchester)
 Hui, L., & Gnedin, N.Y. 1997, *MNRAS*, 292, 27
 Hurley-Keller, D., Mateo, M., & Nemec, J. 1998, *AJ*, 115, 1840
 Ibata, R.A., Gilmore, G., & Irwin, M.J. 1994, *Nature*, 370, 194
 Kepner, J.V., Babul, A., & Spergel, D.N. 1997, *ApJ*, 487, 61
 Kitayama, T., & Ikeuchi, S. 2000, *ApJ*, 529, 615
 Klypin, A., Kravtsov, A.V., Valenzuela, O., & Prada, F. 1999, *ApJ*, 522, 82
 Kormendy, J., & Freeman, K.C. 1998, *AAS*, 193, 2105
 Lin, D.N.C., & Faber, S.M. 1983, *ApJ*, 266, L21
 Mac Low, M., & Ferrara, A. 1999, *ApJ*, 513, 142
 Mateo, M. 1998, *ARAA*, 36, 435
 Miralda-Escudé, J., & Rees, M.J. 1994, *MNRAS*, 266, 343
 Monikiewicz, J., et al. 1999, *PASP*, 111, 1392
 Navarro, J.F., & Steinmetz, M. 1997, *ApJ*, 478, 13
 Ostriker, J.P., & Steinhardt, P. 1995, *Nature*, 377, 600
 Peebles, P.J.E. 1997, *ApJ*, 483, L1
 Press, W.H., & Schechter, P.L. 1974, *ApJ*, 215, 703
 Quinn, T., Katz, N., & Efstathiou, G. 1996, *MNRAS*, 278, L49
 Saviane, I., Held, E.V., & Bertelli, G. 2000, *A&A*, 355, 56
 Schaye, J., Theuns, T., Rauch, M., Efstathiou, G., & Sargent, W.L.W. 2000, *MNRAS*, 318, 817
 Stockman, H.S., & Mather, J. 1999, in "Galaxy Interactions at Low and High Redshift", *Proc. of IAU Symp* 186, ed. J.E. Barnes and D.B. Sanders, p493
 Tegmark, M., et al. 1997, *ApJ*, 474, 1
 Thoul, A.A., & Weinberg, D.H. 1996, *ApJ*, 465, 608
 van den Bergh, S. 2000, "The Galaxies of the Local Group"
 Young, L.M. 2000, *AJ*, 119, 188 (Cambridge University Press)